

DAFTAR PUSTAKA

- Aardema MW. (2001). *Uterine arteri Doppler flow and uteroplacental vascular pathology in normal pregnancies and pregnancies complicated by pre-eclampsia and small for gestational age fetuses*. Br J Obstet Gynaecol.
- Arngrimsson R. (2000). *Genetic and familial predisposition to eclampsia and pre-eclampsia in a defined population*. Br J Obstet Gynaecol.
- Basso, O. (2001). *Higher risk of pre-eclampsia after change of partner. An effect of longer interpregnancy intervals*. Epidemiology.
- Biro Pusat Statistik Indonesia. (2003). *Statistik Indonesia*. Biro Pusat Statistik. Indonesia.
- Catalyst Consortium. (2002) *Optimal Brth Spacing. An Activity of the Catalyst Consortium*. www.rhcatalyst.org.
- Conde-Agudelo, A. (2002). *Maternal morbidity associated with interpregnancy interval: cross sectional study*. BMJ.
- Coonrod, D.V. (1995). *Risk factors for preeclampsia in twin pregnancies: a population based cohort study*. Obstetrics dan Gynecology.
- Dasuki D. (2000). *Preeclampsia-Eclampsia as the single disease and the reproductive risk factors*. Berkala Ilmu Kedokteran.
- Dekker, G.A., Robillard. (2003). *The birth interval hypotesis-does it really indicate the end of the primipaternity hypothesis*. Journal of Reproductive Immunology.
- Dekker, G.A.(2004). *Etiology of Preeclampsia: An Update*. J Med Assoc Thai.
- Dewi Prihastuti. (2004). *Analisis Lanjut SDKI 2002 - 2003, Kecenderungan Preferensi Fertilitas, Unmetneed dan Kehamilan Tidak Diharapkan di Indonesia*. BKKBN, 2004
- Esplin MS.(2001). *Paternal and maternal components of the predisposition to preeclampsia*. N Engl J Med 2001
- Family Care International. (2008). *Family Care International and Safe Motherhood Inter-Agency Group. Safe Motherhood Fact Sheet: 11 fact sheet prepared from the Safe motherhood Technical Consultation in Srilangka*.

- Holland, W.W., (1985). *Oxford Textbook of Public Health: Investigative methods in public health*. Oxford medical publications. Oxford University Press.
- Munro P.T. (2000). *Management of eclampsia in the accident and emergency department*. J. Accid Emerg Med.
- Ness RB (1996). *Heterogeneous causes constituting the single syndrome of preeclampsia: a hypothesis and its implications*. Am J Obstet Gynecol
- Qiu, C. (2003). *Family History of Hypertension and Type 2 Diabetes in Relation to Preeclampsia Risk*. .
- Pipkin, F.B. (2003). *Risk factors for preeclampsia*. New England Journal of Medicine.
- Roberts, J.M., Pearson, G. Cutler, J. dan Linheimer, M. (2003). *Summary of the NHLBI Working Group on Research on Hypertension During Pregnancy*. Hypertension.
- Sibai, B.M.. (1998). *Maternal Perinatal Outcome of Conservative Management of Severe Preeclampsia in mid-trimester*. American Journal Obstetric and Gynecology.
- Sibai, B.M. (2000). *Prevention of preeclampsia: a big disappointment*. American Journal Obstetrics and Gynecology.
- Skjaerven, R.(2002).*The Interval Between Pregnancies and Risk of Preeclampsia*. New England Journal .
- Srinivason, K. (1979). *Birth Interval in Analysis in Fertility Surveys Scientific Report*. Bombay. India
- Staftlas E. (2000). *Epidemiology of preeclampsia and eclampsia in the United States*, Am J Obstet Gynecol.
- Suratman, A.I. (2000). *Evaluasi Penggunaan Dekametason pada Preeklampsia berat dan eklampsia. Kajian Terhadap Morbiditas dan Mortalitas Maternal*. Tesis/Karya Ilmiah Akhir PPDS I Fakultas Kedokteran Universitas Gadjah Mada/RSUP Dr. Sardjito Yogyakarta.
- Trongstad, L.I. (2001). *Changing paternity and time since last pregnancy; the impact on pre-eclampsia risk. A study of 547238 women with and without previous pre-eclampsia*. International Journal of Epidemiology.